



Young Stroke

Defining Young Stroke

Introduction

- | | |
|----------------------------|-------------------|
| 1. Definition | 5. Management |
| 2. Epidemiology | 6. Rehabilitation |
| 3. Risk Factors and Causes | 7. Prevention |
| 4. Diagnosis | 8. Prognosis |

Introduction

Stroke occurs when blood flow to the brain is disturbed, resulting in cell death and dysfunction in one or many parts of the brain. The two main types of stroke include ischemic, due to lack of arterial blood flow to the brain, and hemorrhagic due to bleeding (intracerebral hemorrhage). Although a majority of strokes occur in people aged above 65 years, when it occurs to an adult between the age of 18 and 65 years, it is referred to as a young stroke. A stroke may also occur in children, and its causes may be different from that in young adults and in individuals older than 65 years.

Generally, 80% to 85% of strokes are of ischemic origin and the remaining 15 to 20% are hemorrhagic, but in young stroke a greater proportion are hemorrhagic. A third main type of stroke occurs if venous blood flow from the brain is blocked, leading to cerebral venous thrombosis. This type of stroke generally causes only <1% of all strokes, but it causes about 5% of strokes in the young.

Young stroke should be viewed separately from older strokes for several reasons. The young have a different profile of risk factors and causes underlying their strokes compared with those seen in older stroke patients. Many causes or risk factors more common in young stroke patients are recognized, and include pregnancy and the postpartum period, genetic factors, migraine, illicit drug use, and patent foramen ovale. However, risk factors and causes that are commonly associated with older onset stroke can be present at younger ages as well.

Another important aspect that differentiates the young stroke patients from the older ones is their social and vocational situation. They often have family to take care of, are at the childbearing age, and their strokes often occur at a time of decisive career moves. Although the chances of surviving a stroke at a young age is much better than when it occurs at an older age, they still have to face an increased risk of recurrent strokes for years after the first event. Nevertheless, efficient acute treatment, rehabilitation, and stroke prevention can potentially lead to improved outcomes in individuals with young stroke.

1. Definition

There is no uniform criterion to define the entity “stroke in the young”. In the scientific literature on young adult stroke, the upper age cut-offs have varied from 30 to 65 years, being most commonly 50 or 55 in the recent literature. Although most of the scientific data come from studies using a lower age cut-off, in this article “stroke in the young” refers to ischemic or hemorrhagic strokes occurring in adult people aged less than 65 years, i.e. reflecting usual working age in most parts of the world. This is also the cut-off used in the World Health Organization’s Global Burden of Diseases analyses.

Synonyms to “stroke in the young” include “young-onset stroke”, “early-onset stroke”, and even “young stroke”. There is much less information available on young-onset hemorrhagic than ischemic stroke, and much less information is available on transient ischemic attacks (TIA)—i.e. symptoms originating presumably from a blood clot in an artery that rapidly resolves spontaneously without brain damage seen in imaging studies.

2. Epidemiology

Approximately one fourth of all strokes occur at age of less than 65 years and 1 out of 10 occur at age of less than 50 years.(Ferro, Massaro and Mas, 2010) Incidence is a measure of probability of occurrence of a disease in a population with a specified period of time. Incidence of first stroke increases virtually exponentially with age at younger ages, with the steepest phase of increase beginning in early midlife.(Putala, *et al.*, 2009)

The incidence of ischemic stroke in people aged less than 50 years is about 10 per 100 000 inhabitants per year.(Naess, *et al.*, 2002; Putala, *et al.*, 2009) Among those aged less than 35 years, women outnumber men in the incidence of ischemic stroke, but in those aged more than 35 years but less than 50 years, men outnumber women.(Putala, *et al.*, 2009; Putala, *et al.*, 2012)

Incidence of intracerebral hemorrhage in the young is about a half of that for ischemic stroke. One study showed an approximated incidence of 4.9 per 100.000 for nontraumatic intracerebral hemorrhage at age of 16 to 49 years.(Koivunen, *et al.*, 2014a)

Incidence of cerebral venous thrombosis is unknown, but it encompasses less than 1% of all strokes at all ages. Among younger patients, cerebral venous thrombosis cover up to 5% of all strokes.(Bousser and Ferro, 2007)

There are also notable differences in the incidence between geographic areas and race-ethnic groups. For instance, in the Baltimore-Washington area, incidence of ischemic stroke for young white people was 10.5 per 100.000 while it was 21.7 per 100.000 for young blacks.(Kittner, *et al.*, 1996) As for ischemic stroke, higher rate of intracerebral hemorrhage has been found for young blacks than for young whites.(Kittner, *et al.*, 1996)

Notably, incidence of ischemic stroke in the young has been increasing worldwide since 1980s to present.(Bejot, *et al.*, 2014; Kissela, *et al.*, 2012; Rosengren, *et al.*, 2013; Vangen-Lonne, *et al.*, 2015) The exact reasons for the increase remain unknown, but better awareness of the disease, improved diagnostics, increasing prevalence of overweight and diabetes, unhealthy diet and sedentary lifestyle may play a role. In conjunct with the increasing incidence and decreased stroke death rates, the number of prevalent cases has been increasing globally in the past 20 years. Recent epidemiological findings also suggest that the overall burden of young stroke represents globally now nearly half of the total stroke burden, with most of the burden in developing countries.(Krishnamurthi, *et al.*, 2015)

3. Risk Factors and Causes

Ischemic stroke or TIA

The same risk factors predisposing to ischemic stroke or TIA at older ages may also predispose to young-onset ischemic stroke or TIA. These include hypertension, dyslipidemia, smoking, diabetes mellitus, and cardiovascular disease. In fact, recent studies have shown that such traditional risk factors are common among young stroke patients.(Putaala, *et al.*, 2012; Rolfs, *et al.*, 2013; von Sarnowski, *et al.*, 2013)

There also are specific risk factors to young age. Some risk factors can exist only at young ages (e.g. pregnancy or postpartum period, oral contraceptive use). Other risk factors can exist at all ages, but their influence on the stroke risk has been shown to be stronger at younger ages (e.g. patent foramen ovale, migraine with aura, infections). The factor may also be related to lifestyle-related feature in the population, which is more common at younger ages (e.g. illicit drug use, heavy drinking, binge drinking, smoking, overweight).(Putaala, Heikinheimo-Connell and Tatlisumak. 2015)

The spectrum of causes underlying young-onset ischemic stroke differ markedly from those seen in the elderly patients.(Fromm, *et al.*, 2011) The major causes of ischemic stroke in the elderly, atherosclerosis in the large brain supplying arteries, atrial fibrillation, and small-vessel disease of the brain are rather infrequent among the young. This occurs probably because most traditional risk factors predisposing to these outcomes, such as hypertension, diabetes, and dyslipidemia, are either less common or less severe, or have not yet caused substantial damage to the cardiovascular system.(Ferro, Massaro and Mas, 2010; Yesilot Barlas, *et al.*, 2013)

The most common solitary cause of ischemic stroke in the young in high-income countries is a dissection of a neck artery (a tear within the wall of a blood vessel allows blood to separate the wall layers), involving carotid or vertebral artery. Sometimes a dissection can occur in or extend to an intracranial artery. Dissections alone cause about 15-20% of ischemic stroke in young people. (Debette and Leys, 2009) Other causes seen relatively often among young patients but rarely among elderly patients appear in the table:

- Dissection of a carotid, vertebral, or intracranial artery
- Systemic vasculitides
- Primary angiitis (vasculitis) of the central nervous system
- Systemic lupus erythematosus
- Primary antiphospholipid syndrome
- Sickle cell disease
- Genetic thrombophilic diseases (protein C, protein S, and antithrombin III deficiency; Factor V Leiden and prothrombin gene mutations)
- Sneddon's syndrome
- Moyamoya disease or syndrome
- CADASIL (cerebral autosomal-dominant arteriopathy with subcortical infarcts and leukoencephalopathy)

- Fabry disease
- MELAS (mitochondrial encephalomyopathy, lactic acidosis, and stroke-like episodes)
- HANAC (hereditary angiopathy, nephropathy, aneurysm, and muscle cramps)
- RCVS (reversible cerebral vasoconstriction syndrome)
- Migrainous infarction

Migraine attack resulting in ischemic stroke (migrainous infarction) is a rare occasion with unknown mechanisms, and the diagnosis necessitates exclusion of other potential causes.(Laurell, *et al.*, 2011) Many diseases and conditions caused by a mutation in a single gene may underlie young-onset ischemic stroke, although known mutations are causing each only a fraction of all ischemic strokes in the young.(Cheng, *et al.*, 2014)

Cardiomyopathies, endocarditis, congenital cardiac malformations, atrial fibrillation, endocarditis, and cardiac tumors are examples of high-risk heart conditions that may lead to ischemic stroke at young age. These conditions encompass together about one tenth of all young-onset ischemic strokes.(Yesilot Barlas, *et al.*, 2013) Notably, rheumatic valvular disease of the heart is among the most frequent causes of young-onset ischemic stroke in many non-industrialized countries.(Ghandehari and Moud, 2006)

The reason for the ischemic stroke in the young remains unexplained (i.e. cryptogenic or undetermined) in up to 50% of patients, being more frequent the younger the patient's age.(Yesilot Barlas, *et al.*, 2013) Patent foramen ovale (PFO), a remnant of the fetal circulation that may cause venous blood to enter into arterial side via a shunt between atriums of the heart, is frequently found by cardiac ultrasound examination in these patients.(Kizer and Devereux, 2005) As PFO is also present in about a fourth of healthy individuals, determining when PFO is causally relevant to the ischemic stroke remains a major challenge, however.(Alsheikh-Ali, Thaler and Kent, 2009)

Intracerebral hemorrhage

As in older stroke patients, hypertension and smoking are the most frequent risk factors for young-onset hemorrhagic stroke. However, among the young, illicit drug use, hematologic disorders, and genetic conditions may be predisposing to hemorrhagic stroke in the young more often than in older patients.(Koivunen, *et al.*, 2014a)

As for ischemic stroke, the spectrum of causes for hemorrhagic stroke at younger ages is variable and different from that in the elderly. Arteriopathy related to hypertension is less frequent and amyloid angiopathy absent, while vascular malformations, hematologic disorders, illicit drug use, moyamoya disease, and vasculitides are more commonly causing intracerebral hemorrhage in young adults. Also for intracerebral hemorrhage, the cause remains unknown in relatively many young patient.(Koivunen, *et al.*, 2014a)

Causes atypical in elderly but relatively often underlying hemorrhagic stroke among young adults include:

- Arteriovenous malformation
- Cavernoma
- Coagulopathies and hematologic disorders
- Systemic vasculitis or primary angiitis (vasculitis) of the central nervous system
- RCVS (reversible cerebral vasoconstriction syndrome)

- Illicit drug use
- Moyamoya disease
- Cerebral venous thrombosis
- Pre-eclampsia or eclampsia

Cerebral venous thrombosis

Risk factors and reasons for cerebral venous thrombosis include pregnancy and puerperium, infections, systemic inflammatory conditions, genetic prothrombotic conditions, hematologic conditions, certain drugs (e.g. oral contraceptives, hormone replacement therapy, steroids), mechanical causes and trauma, as well as dehydration and cancer. Cerebral venous thrombosis can cause both ischemic or hemorrhagic lesion in the brain.(Bousser and Ferro, 2007)

4. Diagnosis

The basic diagnostic work-up does not differ in young stroke patients from that in older patients. However, due to the extensive array of different causes, a vigorous interview of patient and family history, clinical examination, and specific ancillary tests are often needed besides routine brain and vascular imaging, laboratory tests, and cardiac ultrasound.

5. Management

General acute management of young-onset stroke is similar to that in older patients. Acute treatment strategies of ischemic stroke include immediate transport to a hospital providing acute stroke care, administration of intravenous thrombolysis with or without endovascular thrombectomy within the first hours after symptom onset, followed by treatment in a stroke unit.(Jauch, *et al.*, 2013; Powers, *et al.*, 2015) Younger patients may benefit from intravenous thrombolysis with greater extent compared with older patients with similar symptom severity.(Toni, *et al.*, 2012)

In very large middle cerebral artery infarctions, a neurosurgical procedure called hemicraniectomy may be lifesaving, and, in the best scenario, affect favorably to functional outcome (i.e. ability to move, independency in daily living). In this procedure, part of the skull is removed to allow a swelling brain to expand without being squeezed to the opposite side of the brain.(Vahedi, *et al.*, 2007) Also for hemicraniectomy, younger patients may have more benefit from the procedure than older patients.(Juttler, *et al.*, 2014; Vahedi, *et al.*, 2007)

Treatment of intracerebral hemorrhage also follows the same principles as for older patients.(Hemphill, *et al.*, 2015) Because vascular malformations more often underlie intracerebral hemorrhage in the young, surgical or endovascular treatments may be more often warranted.(Koivunen, *et al.*, 2014b)

Cerebral venous thrombosis is usually treated with oral anticoagulants for 3 to 6 months. Depending on the underlying cause, the need for anticoagulation treatment may occasionally be permanent.(Bousser and Ferro, 2007)

6. Rehabilitation

Individuals with young stroke have unique rehabilitation needs. Both physical and cognitive skills are impacted by stroke, which has profound effects on a young individual's quality of life.(Keppel & Crowe 2000; O'Connor, *et al.*, 2005; Röding, *et al.*, 2009) This is because they are at a stage in life when employment is important, and are more likely to be caring for children.(Dixon, *et al.*, 2007; Stone 2007) However, few studies have assessed the specific needs of young stroke patients.(Low, *et al.*, 2003) As a result, many of the needs of young stroke patients are unmet in the context of inpatient rehabilitation which tends to focus on older stroke patients.(Röding, *et al.*, 2003; Stone 2005)

Furthermore, young stroke survivors have a longer period of disability to contend with due to their impairments which creates a socioeconomic challenge.(Bjorkdahl & Sunnerhagen 2007; Mehndiratta, *et al.*, 2004, O'Connor, *et al.*, 2005) The cost of stroke in young people exceeds that of stroke in older people due to a loss in productivity and more psychosocial complications.(Jacobs, *et al.*, 2002; Nayak, *et al.*, 1997) These complications include stress in the family, institutionalization, return to work and future needs.(Teasell, *et al.*, 2000) Thus the rehabilitation of young stroke patients presents distinct rehabilitation challenges.

Young stroke patients have been shown to demonstrate greater neurological and functional recovery and have a better prognosis and long-term survival rate compared to older stroke patients.(Adunsky, *et al.*, 1992; Hindfelt & Nilsson 1992; Marini, *et al.*, 2001; Nedeltchev, *et al.*, 2005, Black-Schaffer & Winston 2004) However, they are at increased risk for experiencing reduced cognitive performance, anxiety and depression compared to stroke-free age and sex matched controls.(Schaapsmeeders, *et al.*, 2013, Waje-Andreassen, *et al.*, 2013)

Traditional rehabilitation is generally the same for younger and older stroke patients.(Teasell, *et al.*, 2000) Rehabilitation starts in the acute care hospital, and continues on inpatient rehabilitation units, then in outpatient therapy programs, and eventually progress to vocational rehabilitation. Following stroke there can be a period of natural recovery which is characterized by a non-linear pattern of improvement. This can look like a stairstep pattern of improvement across different areas. It is now well-established that recovery may continue for many years after the stroke.(Sabini, *et al.*, 2012) Specialty guidelines for stroke rehabilitation have been developed (www.carf.org, and www.strokebestpractices.ca/). Return to work or to other life activities such as school, or parenting are important goals of rehabilitation. The unique issues associated with younger stroke rehabilitation pertain to the nature of family support, the presence of young dependents, marital stress, and return to work.(Dixon, *et al.* 2007)

Cognitive impairment from stroke can complicate a young stroke patient's reintegration back into the community and return to work.(Malm, *et al.*, 1998, Hommel, *et al.*, 2009) Almost half of young stroke patients are diagnosed with some degree of post-stroke depression ,(Kappelle, *et al.*, 1994; Neau, *et al.*, 1998) which is associated with a poor rehabilitation outcome (Neau, *et al.*, 1998). A better understanding of cognitive deficits is required for more effective age-adapted rehabilitation programs.(Röding, *et al.*, 2003) Young stroke patients also expressed a need for communication with other stroke patients their age that had been affected by similar experiences.(Dixon, *et al.*, 2007; Röding, *et al.* 2003) Furthermore, rehabilitation for young stroke survivors should emphasize participation in fitness activities, as they are necessary for well-being and secondary stroke prevention ;(Naess, *et al.*, 2004) unfortunately these activities are often given up following a stroke. (Wolf, *et al.*, 2012) Post stroke fatigue in young adults can negatively affect scholastic, vocational, and social pursuits. It is associated with higher mortality rates, depression, and lower functional outcomes in older patients,(Naess, *et al.*, 2005) but is relatively understudied in young survivors.

Individuals interested in rehabilitation services should seek consultation from a neurologist or rehabilitation physician specializing in stroke rehabilitation. Due to the interaction between physical symptoms, changes in thinking skills, and emotional changes, persons with stroke can benefit from broad-based specialty rehabilitation therapies by an interdisciplinary team, including physical, occupational, speech, recreational therapists and specialists in physical medicine and neuropsychology. Common rehabilitation regimes include some combination of physiotherapy, speech language therapy, occupational therapy, and pharmacological therapy. (Stein 2004; Young & Forster 2007) Strategies to improve motor recovery in young stroke patients include: Constraint-induced movement therapy, robot-aided rehabilitation, virtual reality training, EMG-biofeedback, functional electrical stimulation, increased exercise intensity, and acupuncture.(Stein 2004)

A stroke affects the entire family. Caregivers of stroke survivors suffer from higher rates of depression and greater rates of deterioration in their own health.(Kinsella & Duffy 1979) Caregivers report more emotional distress when caring for individuals with stroke who exhibit more depressive symptoms and cognitive impairment, and when caregivers were younger, female, in poorer physical health, and experienced more interference with their lifestyle. (Cameron, *et al.*, 2011) In the case of younger strokes, where the primary caregiver is the spouse, added responsibilities of caring for children create further adjustment problems.(Visser-Meily, *et al.*, 2005, Martinsen, *et al.*, 2012) The relationship between parents post-stroke and their children is complex, and the stroke event can add stress to the family environment.

Vocational and return to work issues are a major facet of rehabilitation unique to the young stroke population.(Malm, *et al.*, 1998) Significant predictors of return to work include age, sex, functional status, absence of psychiatric illness, and education level.(Bergmann, *et al.*, 1991; Glozier, *et al.*, 2008; Howard, *et al.*, 1985; Peters, *et al.*, 2013) Individuals in managerial positions and white collar occupations are most likely to return to work.(Howard 1995, Saeki, *et al.*, 1993; 2010) The higher success rate for white collar positions may be attributed to better education, work conditions, pay and less physically demanding tasks. The side of hemiplegia or weakness has not been consistently associated with the ability to return to work in young stroke patients.(Black-Schaffer & Osberg 1990; Heinemann, *et al.*, 1987; Howard, *et al.*, 1985; Kotila, *et al.* 1984; Weisbroth, *et al.*, 1971) However, those with better upper extremity use, ambulation, and abstract reasoning are more likely to return to work. Right hemiplegics with milder communication and cognitive deficits also have better vocational outcomes. There is a distinct negative correlation between aphasia and return to work.(Black-Schaffer & Osberg 1990) Neuropsychological testing may be required in order to accurately delineate the extent of cognitive problems and determine how they might impact the patient's eventual return to work.(Lindberg, *et al.*, 1992; Ljunggren, *et al.*, 1985). The inability to return to work frequently leads to emotional and financial hardships for stroke survivors and their families.(Churchill 1993) To address this issue, The Stroke Association & Different Strokes published *Getting Back to Work after Stroke*.(Barker 2006) Employers and professionals within the community should try to focus on meeting the needs of stroke survivors and their families in order to facilitate participation and independence for stroke patients.(Kersten, *et al.*, 2002) Young stroke patients need to be connected with support organizations and those who share similar experiences (see www.youngstroke.org).

7. Prevention

Primary prevention of stroke at all ages includes generally healthy lifestyle, in particular regular physical activity, weight control, healthy diet favoring Mediterranean diet ingredients, and avoidance of tobacco smoking, excess alcohol use and illicit drugs.(Meschia, *et al.*, 2014)

Secondary prevention after a stroke or TIA at young age should generally follow the same principles as for older patients.(Kernan, *et al.*, 2014) Medication and preventive measures

should be tailored according to the underlying cause and therefore it is of utmost importance to carry out extensive and timely diagnostic investigations to decipher the cause for the stroke in a young patient.

If established vascular risk factors (e.g. elevated blood pressure, high cholesterol, diabetes mellitus) are present, their careful treatment is warranted. Medical treatment typically involves antiplatelet medication or anticoagulants. Depending on clinical features, these are combined with cholesterol and/or blood pressure lowering medications. If a specific cause—such as stenosis in a cervical artery due to atherosclerosis, cardiac myxoma, arteriovenous malformation, moyamoya disease—is present, the preventive treatment aims at removing such a source or creating a bypass in the brain circulation. When a systemic condition such as a hematologic disease or systemic vasculitis is primarily underlying the stroke, its efficient treatment forms a cornerstone of secondary prevention. Estrogen-containing hormonal product should be stopped. Other precipitants such as excess alcohol drinking, illicit drug use should be eliminated to minimize the risk of future strokes.

In many conditions related to young-onset stroke, there is uncertainty regarding the optimal secondary prevention. Patent foramen ovale is an example of such condition and currently it remains uncertain whether closure of patent foramen ovale, using a catheter device, is superior to best medical treatment.(Li, *et al.*, 2015) It also remains uncertain for many cases, how long secondary preventive medication should be continued, particularly if there are no known vascular risk factors present and the stroke remained cryptogenic.(Naess, *et al.*, 2005)

8. Prognosis

Age is the most important factor driving mortality after stroke. Young stroke patients thus generally survive their strokes more often compared with older patients. However, compared with background population of same age, young stroke patients face an about 4-fold risk of dying in the long-term after their stroke.(Rutten-Jacobs, *et al.*, 2013a) The most frequent causes of death include vascular diseases,(Putala, *et al.*, 2009) so their efficient treatment is highly important to mitigate the risk of dying after young-onset stroke.

Observational studies show that young patients with ischemic stroke have a substantial risk of future nonfatal strokes and other cardiovascular events, depending primarily on the risk factors and cause of the initial stroke.(Pezzini, *et al.*, 2014; Putala, *et al.*, 2010; Rutten-Jacobs, *et al.*, 2013b)

Functional outcome in young stroke patients is generally favorable, with about 4 out of 5 regaining independence in daily living measured with modified Rankin Scale.(Naess, *et al.*, 2002; Putala, *et al.*, 2013) However, such scales are only crude measures of clinical outcome and they may not accurately consider for example subtle cognitive deficits, fatigue, and depression, which are rather common residual symptoms in young patients after stroke, and may hamper quality of life.(Koivunen, *et al.*, 2015; Maaijwee, *et al.*, 2015; Naess, *et al.*, 2006; Synhaeve, *et al.*, 2015)

Return to work and productive duties is among the most important outcome after young-onset stroke as mentioned above. Limited number of studies including patients from many decades indicates that rather high percentage of young stroke survivors is left without paid work in the long term.(Maaijwee, *et al.*, 2014) Work adjustments may also be needed for those returning to work.

References

- Adunsky, A., Hershkowitz, M., Rabbi, R., Asher-Sivron, L., & Ohry, A. (1992). Functional recovery in young stroke patients. *Arch.Phys.Med.Rehabil.*, 73(9), 859-862.
- Alsheikh-Ali AA, Thaler DE, Kent DM. Patent foramen ovale in cryptogenic stroke: Incidental or pathogenic? *Stroke* 2009; 40: 2349-55.
- Barker, G. (2006). *Getting back to work after stroke*. London: The Stroke Association & Different Stroke, United Kingdom.
- Bejot Y, Daubail B, Jacquin A, Durier J, Osseby GV, Rouaud O, et al. Trends in the incidence of ischaemic stroke in young adults between 1985 and 2011: The dijon stroke registry. *J Neurol Neurosurg Psychiatry* 2014; 85: 509-13.
- Bergmann, H., Kuthmann, M., von Ungern-Sternberg, A., & Weimann, V. G. (1991). Medical educational and functional determinants of employment after stroke. *J.Neural Transm.Suppl*, 33, 157-161.
- Bjorkdahl, A., & Sunnerhagen, K. S. (2007). Process skill rather than motor skill seems to be a predictor of costs for rehabilitation after a stroke in working age; a longitudinal study with a 1 year follow up post discharge. *BMC.Health Serv.Res.*, 7, 209.
- Black-Schaffer, R. M., & Osberg, J. S. (1990). Return to work after stroke: development of a predictive model. *Arch.Phys.Med.Rehabil.*, 71(5), 285-290.
- Black-Schaffer, R. M., & Winston, C. (2004). Age and functional outcome after stroke. *Top.Stroke Rehabil.*, 11(2), 23-32.
- Bousser MG, Ferro JM. Cerebral venous thrombosis: An update. *Lancet Neurol* 2007; 6: 162-70.
- Cameron, J. I., Cheung, A. M., Streiner, D. L., Coyte, P. C., & Stewart, D. E. (2011). Stroke survivor depressive symptoms are associated with family caregiver depression during the first 2 years poststroke. *Stroke*, 42(2), 302-306.
- Cheng YC, Cole JW, Kittner SJ, Mitchell BD. Genetics of ischemic stroke in young adults. *Circ Cardiovasc Genet* 2014; 7: 383-92.
- Churchill, C. (1993). Social Problems Post stroke. In R. W. Teasell (Ed.), *Phys Med Rehab: State of the Art Reviews*. Long term consequences of stroke (Vol. 7, pp. 213-223). Philadelphia: Hanley & Belfus.
- Debette S, Leys D. Cervical-artery dissections: Predisposing factors, diagnosis, and outcome. *Lancet Neurol* 2009; 8: 668-78.
- Dixon, G., Thornton, E. W., & Young, C. A. (2007). Perceptions of self-efficacy and rehabilitation among neurologically disabled adults. *Clin.Rehabil.*, 21(3), 230-240.
- Ferro JM, Massaro AR, Mas JL. Aetiological diagnosis of ischaemic stroke in young adults. *Lancet Neurol* 2010; 9: 1085-96.

Fromm A, Waje-Andreassen U, Thomassen L, Naess H. Comparison between ischemic stroke patients ≥ 50 years admitted to a single centre: The bergen stroke study. *Stroke Res Treat* 2011; 2011: 183256.

Ghandehari K, Moud ZI. Incidence and etiology of ischemic stroke in persian young adults. *Acta Neurol Scand* 2006; 113: 121-4.

Glozier, N., Hackett, M. L., Parag, V., & Anderson, C. S. (2008). The influence of psychiatric morbidity on return to paid work after stroke in younger adults: the Auckland Regional Community Stroke (ARCOS) Study, 2002 to 2003. *Stroke*, 39(5), 1526-1532.

Heinemann, A. W., Roth, E. J., Cichowski, K., & Betts, H. B. (1987). Multivariate analysis of improvement and outcome following stroke rehabilitation. *Archives of neurology*, 44(11), 1167.

Hemphill JC, 3rd, Greenberg SM, Anderson CS, Becker K, Bendok BR, Cushman M, et al. Guidelines for the management of spontaneous intracerebral hemorrhage: A guideline for healthcare professionals from the american heart Association/American stroke association. *Stroke* 2015; 46: 2032-60.

Hindfelt, B., & Nilsson, O. (1992). Long-term prognosis of ischemic stroke in young adults. *Acta Neurol.Scand.*, 86(5), 440-445.

Hommel, M., Miguel, S. T., Naegele, B., Gonnet, N., & Jaillard, A. (2009). Cognitive determinants of social functioning after a first ever mild to moderate stroke at vocational age. *J.Neurol.Neurosurg.Psychiatry*, 80(8), 876-880.

Howard, G., Till, J. S., Toole, J. F., Matthews, C., & Truscott, B. L. (1985). Factors influencing return to work following cerebral infarction. *JAMA: the journal of the American Medical Association*, 253(2), 226-232.

Jacobs, B. S., Boden-Albala, B., Lin, I. F., & Sacco, R. L. (2002). Stroke in the young in the northern Manhattan stroke study. *Stroke*, 33(12), 2789-2793.

Jauch EC, Saver JL, Adams HP, Jr, Bruno A, Connors JJ, Demaerschalk BM, et al. Guidelines for the early management of patients with acute ischemic stroke: A guideline for healthcare professionals from the american heart Association/American stroke association. *Stroke* 2013; 44: 870-947.

Juttler E, Unterberg A, Woitzik J, Bosel J, Amiri H, Sakowitz OW, et al. Hemispherectomy in older patients with extensive middle-cerebral-artery stroke. *N Engl J Med* 2014; 370: 1091-100.

Kappelle, L. J., Adams, H. P., Jr., Heffner, M. L., Torner, J. C., Gomez, F., & Biller, J. (1994). Prognosis of young adults with ischemic stroke. A long-term follow-up study assessing recurrent vascular events and functional outcome in the Iowa Registry of Stroke in Young Adults. *Stroke*, 25(7), 1360-1365.

Keppel, C. C., & Crowe, S. F. (2000). Changes to body image and self-esteem following stroke in young adults. *Neuropsychological Rehabilitation*, 10(1), 15-31.

Kernan WN, Ovbiagele B, Black HR, Bravata DM, Chimowitz MI, Ezekowitz MD, et al. Guidelines for the prevention of stroke in patients with stroke and transient

ischemic attack: A guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2014; 45: 2160-236.

Kersten, P., Low, J. T., Ashburn, A., George, S. L., & McLellan, D. L. (2002). The unmet needs of young people who have had a stroke: results of a national UK survey. *Disabil.Rehabil.*, 24(16), 860-866.

Kinsella, G. J., & Duffy, F. D. (1979). Psychosocial readjustment in the spouses of aphasic patients. A comparative Survey of 79 subjects. *Scand.J.Rehabil.Med.*, 11(3), 129-132.

Kissela BM, Khoury JC, Alwell K, Moomaw CJ, Woo D, Adeoye O, et al. Age at stroke: Temporal trends in stroke incidence in a large, biracial population. *Neurology* 2012; 79: 1781-7.

Kittner SJ, Stern BJ, Feaser BR, Hebel R, Nagey DA, Buchholz DW, et al. Pregnancy and the risk of stroke. *N Engl J Med* 1996; 335: 768-74.

Kizer JR, Devereux RB. Clinical practice. patent foramen ovale in young adults with unexplained stroke. *N Engl J Med* 2005; 353: 2361-72.

Koivunen RJ, Harno H, Tattlisumak T, Putaala J. Depression, anxiety, and cognitive functioning after intracerebral hemorrhage. *Acta Neurol Scand* 2015; 132: 179-84.

Koivunen RJ, Satopaa J, Haapaniemi E, Strbian D, Meretoja A, Mustanoja S, et al. Predictors of early mortality in young adults after intracerebral hemorrhage. *Stroke* 2014b; 45: 2454-6.

Koivunen RJ, Satopaa J, Meretoja A, Strbian D, Haapaniemi E, Niemela M, et al. Incidence, risk factors, etiology, severity and short-term outcome of non-traumatic intracerebral hemorrhage in young adults. *Eur J Neurol* 2014a.

Kotila, M., Waltimo, O., Niemi, M. L., Laaksonen, R., & Lempinen, M. (1984). The profile of recovery from stroke and factors influencing outcome. *Stroke*, 15(6), 1039-1044.

Krishnamurthi RV, Moran AE, Feigin VL, Barker-Collo S, Norrving B, Mensah GA, et al. Stroke prevalence, mortality and disability-adjusted life years in adults aged 20-64 years in 1990-2013: Data from the global burden of disease 2013 study. *Neuroepidemiology* 2015; 45: 190-202.

Laurell K, Artto V, Bendtsen L, Hagen K, Kallela M, Meyer EL, et al. Migrainous infarction: A nordic multicenter study. *Eur J Neurol* 2011; 18: 1220-6.

Li J, Liu J, Liu M, Zhang S, Hao Z, Zhang J, et al. Closure versus medical therapy for preventing recurrent stroke in patients with patent foramen ovale and a history of cryptogenic stroke or transient ischemic attack. *Cochrane Database Syst Rev* 2015; 9: CD009938.

Lindberg, M., Angquist, K. A., Fodstad, H., Fugl-Meyer, K., & Fugl-Meyer, A. R. (1992). Self-reported prevalence of disability after subarachnoid haemorrhage, with special emphasis on return to leisure and work. *Br.J.Neurosurg.*, 6(4), 297-304.

Ljunggren, B., Sonesson, B., Saveland, H., & Brandt, L. (1985). Cognitive impairment and adjustment in patients without neurological deficits after aneurysmal SAH and early operation. *J.Neurosurg.*, 62(5), 673-679.

Low, J. T., Kersten, P., Ashburn, A., George, S., & McLellan, D. L. (2003). A study to evaluate the met and unmet needs of members belonging to Young Stroke groups affiliated with the Stroke Association. *Disabil.Rehabil.*, 25(18), 1052-1056.

Maaijwee NA, Arntz RM, Rutten-Jacobs LC, Schaapsmeeders P, Schoonderwaldt HC, van Dijk EJ, et al. Post-stroke fatigue and its association with poor functional outcome after stroke in young adults. *J Neurol Neurosurg Psychiatry* 2015; 86: 1120-6.

Maaijwee NA, Rutten-Jacobs LC, Arntz RM, Schaapsmeeders P, Schoonderwaldt HC, van Dijk EJ, et al. Long-term increased risk of unemployment after young stroke: A long-term follow-up study. *Neurology* 2014; 83: 1132-8.

Malm, J., Kristensen, B., Karlsson, T., Carlberg, B., Fagerlund, M., & Olsson, T. (1998). Cognitive impairment in young adults with infratentorial infarcts. *Neurology*, 51(2), 433-440.

Marini, C., Totaro, R., De, S. F., Ciancarelli, I., Baldassarre, M., & Carolei, A. (2001). Stroke in young adults in the community-based L'Aquila registry: incidence and prognosis. *Stroke*, 32(1), 52-56.

Martinsen, R., Kirkevold, M., & Sveen, U. (2012). Younger stroke survivors' experiences of family life in a long-term perspective: a narrative hermeneutic phenomenological study. *Nurs Res Pract*, 2012, 948791.

Mehndiratta, M. M., Agarwal, P., Sen, K., & Sharma, B. (2004). Stroke in young adults: a study from a university hospital in north India. *Medical science monitor: international medical journal of experimental and clinical research*, 10(9), CR535.

Meschia JF, Bushnell C, Boden-Albala B, Braun LT, Bravata DM, Chaturvedi S, et al. Guidelines for the primary prevention of stroke: A statement for healthcare professionals from the American Heart Association/American stroke association. *Stroke* 2014; 45: 3754-832.

Naess H, Nyland HI, Thomassen L, Aarseth J, Nyland G, Myhr KM. Incidence and short-term outcome of cerebral infarction in young adults in western Norway. *Stroke* 2002; 33: 2105-8.

Naess H, Waje-Andreassen U, Thomassen L, Nyland H, Myhr KM. Do all young ischemic stroke patients need long-term secondary preventive medication? *Neurology* 2005; 65: 609-11.

Naess H, Waje-Andreassen U, Thomassen L, Nyland H, Myhr KM. Health-related quality of life among young adults with ischemic stroke on long-term follow-up. *Stroke* 2006; 37: 1232-6.

Naess, H., Nyland, H. I., Thomassen, L., Aarseth, J., & Myhr, K. M. (2004). Long-term outcome of cerebral infarction in young adults. *Acta Neurol.Scand.*, 110(2), 107-112.

Naess, H., Nyland, H. I., Thomassen, L., Aarseth, J., & Myhr, K. M. (2005). Fatigue at long-term follow-up in young adults with cerebral infarction. *Cerebrovasc.Dis.*, 20(4), 245-250.

Nayak, S. D., Nair, M., Radhakrishnan, K., & Sarma, P. S. (1997). Ischaemic stroke in the young adult: clinical features, risk factors and outcome. *Natl.Med.J.India*, 10(3), 107-112.

Neau, J. P., Ingrand, P., Mouille-Brachet, C., Rosier, M. P., Couderq, C., Alvarez, A., & Gil, R. (1998). Functional recovery and social outcome after cerebral infarction in young adults. *Cerebrovasc.Dis.*, 8(5), 296-302.

Nedeltchev, K., der Maur, T. A., Georgiadis, D., Arnold, M., Caso, V., Mattle, H. P., Schroth, G., Remonda, L., Sturzenegger, M., Fischer, U., & Baumgartner, R. W. (2005). Ischaemic stroke in young adults: predictors of outcome and recurrence. *J.Neurol.Neurosurg.Psychiatry*, 76(2), 191-195.

O'Connor, R. J., Cassidy, E. M., & Delargy, M. A. (2005). Late multidisciplinary rehabilitation in young people after stroke. *Disabil.Rehabil.*, 27(3), 111-116.

Peters, G., Buni, S., Oyeyemi, A., & Hamzat, T. (2013). Determinants of return to work among Nigerian stroke survivors. *Disability & Rehabilitation*, 35(6), 455-459.

Pezzini A, Grassi M, Lodigiani C, Patella R, Gandolfo C, Zini A, et al. Predictors of long-term recurrent vascular events after ischemic stroke at young age: The italian project on stroke in young adults. *Circulation* 2014; 129: 1668-76.

Powers WJ, Derdeyn CP, Biller J, Coffey CS, Hoh BL, Jauch EC, et al. 2015 american heart Association/American stroke association focused update of the 2013 guidelines for the early management of patients with acute ischemic stroke regarding endovascular treatment: A guideline for healthcare professionals from the american heart Association/American stroke association. *Stroke* 2015; 46: 3020-35.

Putala J, Curtze S, Hiltunen S, Tolppanen H, Kaste M, Tatlisumak T. Causes of death and predictors of 5-year mortality in young adults after first-ever ischemic stroke: The helsinki young stroke registry. *Stroke* 2009; 40: 2698-703.

Putala J, Haapaniemi E, Metso AJ, Metso TM, Artto V, Kaste M, et al. Recurrent ischemic events in young adults after first-ever ischemic stroke. *Ann Neurol* 2010; 68: 661-71.

Putala J, Heikinheimo-Connell T, Tatlisumak T. Risk factors specific to young adults. In: Seshadri S, Debette S, editors. *Risk Factors for Cerebrovascular Disease and Stroke*. : Oxford University Press; 2015.

Putala J, Metso AJ, Metso TM, Konkola N, Kraemer Y, Haapaniemi E, et al. Analysis of 1008 consecutive patients aged 15 to 49 with first-ever ischemic stroke: The helsinki young stroke registry. *Stroke* 2009; 40: 1195-203.

Putala J, Strbian D, Mustanoja S, Haapaniemi E, Kaste M, Tatlisumak T. Functional outcome in young adult ischemic stroke: Impact of lipoproteins. *Acta Neurol Scand* 2013; 127: 61-9.

Putala J, Yesilot N, Waje-Andreassen U, Pitkaniemi J, Vassilopoulou S, Nardi K, et al. Demographic and geographic vascular risk factor differences in European young adults with ischemic stroke: The 15 cities young stroke study. *Stroke* 2012; 43: 2624-30.

Rolfs A, Fazekas F, Grittner U, Dichgans M, Martus P, Holzhausen M, et al. Acute cerebrovascular disease in the young: The stroke in young Fabry patients study. *Stroke* 2013; 44: 340-9.

Rosengren A, Jiang KW, Lappas G, Jern C, Toren K, Bjorck L. Twenty-four-year trends in the incidence of ischemic stroke in Sweden from 1987 to 2010. *Stroke* 2013; 44: 2388-93.

Rutten-Jacobs LC, Arntz RM, Maaijwee NA, Schoonderwaldt HC, Dorresteijn LD, van Dijk EJ, et al. Long-term mortality after stroke among adults aged 18 to 50 years. *JAMA* 2013a; 309: 1136-44.

Rutten-Jacobs LC, Maaijwee NA, Arntz RM, Schoonderwaldt HC, Dorresteijn LD, van der Vlugt MJ, et al. Long-term risk of recurrent vascular events after young stroke: The FUTURE study. *Ann Neurol* 2013b; 74: 592-601.

Röding, J. (2009). Stroke in the younger: Self-reported impact on work situation, cognitive function, physical function and life satisfaction-A national survey. Umeå University.

Röding, J., Lindstrom, B., Malm, J., & Ohman, A. (2003). Frustrated and invisible--younger stroke patients' experiences of the rehabilitation process. *Disabil.Rehabil.*, 25(15), 867-874.

Sabini R, Dijkers M, Raghavan P. (2012) Stroke survivors talk while doing: Development of a new therapeutic framework for recovery of hand function post stroke. *Journal of Hand Therapy*; 26(2):124-30.

Saeki, S., & Toyonaga, T. (2010). Determinants of early return to work after first stroke in Japan. *J.Rehabil.Med.*, 42(3), 254-258.

Saeki, S., Ogata, H., Okubo, T., Takahashi, K., & Hoshuyama, T. (1993). Factors influencing return to work after stroke in Japan. *Stroke*, 24(8), 1182-1185.

Schaapsmeeders, P., Maaijwee, N. A., Van Dijk, E. J., Rutten-Jacobs, L. C., Arntz, R. M., Schoonderwaldt, H. C., Dorresteijn, L. D., Kessels, R. P., & de Leeuw, F. E. (2013). Long-term cognitive impairment after first-ever ischemic stroke in young adults. *Stroke*, 44(6), 1621-1628.

Stein, J. (2004). Motor recovery strategies after stroke. *Top.Stroke Rehabil.*, 11(2), 12-22.

Stone, S. D. (2005). Reactions to invisible disability: the experiences of young women survivors of hemorrhagic stroke. *Disabil.Rehabil.*, 27(6), 293-304.

Stone, S. D. (2007). Patient concerns posthaemorrhagic stroke: a study of the Internet narratives of patients with ruptured arteriovenous malformation. *J.Clin.Nurs.*, 16(2), 289-297.

Synhaeve NE, Schaapsmeeders P, Arntz RM, Maaijwee NA, Rutten-Jacobs LC, Schoonderwaldt HC, et al. Cognitive performance and poor long-term functional outcome after young stroke. *Neurology* 2015; 85: 776-82.

Teasell, R. W., McRae, R., & Finestone, H. M. (2000). Social issues in the rehabilitation of younger stroke patients. *Archives of physical medicine and rehabilitation*, 81(2), 205-209.

Toni D, Ahmed N, Anzini A, Lorenzano S, Brozman M, Kaste M, et al. Intravenous thrombolysis in young stroke patients: Results from the SITS-ISTR. *Neurology* 2012; 78: 880-7.

Vahedi K, Hofmeijer J, Juettler E, Vicaut E, George B, Algra A, et al. Early decompressive surgery in malignant infarction of the middle cerebral artery: A pooled analysis of three randomised controlled trials. *Lancet Neurol* 2007; 6: 215-22.

Waje-Andreassen, U., Thomassen, L., Jusufovic, M., Power, K. N., Eide, G. E., Vedeler, C. A., & Naess, H. (2013). Ischaemic stroke at a young age is a serious event - final results of a population-based long-term follow-up in Western Norway. *European Journal of Neurology*, 20(5), 818-823.

Vangen-Lonne AM, Wilsgaard T, Johnsen SH, Carlsson M, Mathiesen EB. Time trends in incidence and case fatality of ischemic stroke: The tromso study 1977-2010. *Stroke* 2015; 46: 1173-9.

Weisbroth, S., Esibill, N., & Zuger, R. R. (1971). Factors in the vocational success of hemiplegic patients. *Archives of physical medicine and rehabilitation*, 52(10), 441.

Visser-Meily, A., Post, M., Meijer, A. M., van, d. P., I, Maas, C., & Lindeman, E. (2005). When a parent has a stroke: clinical course and prediction of mood, behavior problems, and health status of their young children. *Stroke*, 36(11), 2436-2440.

Wolf, T. J., Brey, J. K., Baum, C., & Connor, L. T. (2012). Activity participation differences between younger and older individuals with stroke. *Brain Impairment*, 13(1), 16-23.

von Sarnowski B, Putaala J, Grittner U, Gaertner B, Schminke U, Curtze S, et al. Lifestyle risk factors for ischemic stroke and transient ischemic attack in young adults in the stroke in young fabry patients study. *Stroke* 2013; 44: 119-25.

Yesilot Barlas N, Putaala J, Waje-Andreassen U, Vassilopoulou S, Nardi K, Odier C, et al. Etiology of first-ever ischaemic stroke in european young adults: The 15 cities young stroke study. *Eur J Neurol* 2013; 20: 1431-9.

Young, J., & Forster, A. (2007). Review of stroke rehabilitation. *BMJ*, 334(7584), 86-90.